

Jurnal Info Kesehatan

Vol. 21, No. 4, December 2023, pp. 886-895

P-ISSN 0216-504X, E-ISSN 2620-536X

DOI: [10.31965/infokes.Vol21Iss4.1183](https://doi.org/10.31965/infokes.Vol21Iss4.1183)

Journal homepage: <http://jurnal.poltekkeskupang.ac.id/index.php/infokes>



RESEARCH

Open Access

Pesticide Exposure and Increased Liver Enzyme Activity among Suburban Horticultural

Ahmad Dahlan^{1a}, James Perdinan Simanjuntak^{1b*}, Raden Mustopa^{1c}, Devi Oktarina Putri^{1d}, Putrilia Amanda^{1e}, Adinda Cahyaning Ratri^{1f}, Ahmad Syathibi^{1g}, Sabarudin^{1h}, Haflin¹ⁱ

¹ Department of Environmental Health, Poltekkes Kemenkes Jambi, Jambi City, Jambi Province, Indonesia

^a Email address: dahlan1961@poltekkesjambi.ac.id

^b Email address: james.p.simanjuntak@poltekkesjambi.ac.id

^c Email address: mustopa.rm@poltekkesjambi.ac.id

^d Email address: devioktarinap@gmail.com

^e Email address: putriliaamanda25@gmail.com

^f Email address: adindacahyaningratri@gmail.com

^g Email address: syarthibiahmad@gmail.com

^h Email address: sabarudin.ahmad@gmail.com

ⁱ Email address: drshaflinapt@gmail.com

Received: 14 May 2023

Revised: 23 September 2023

Accepted: 24 September 2023

Abstract

Horticultural farmers use chemicals such as pesticides to increase productivity and also product quality. Exposure to pesticides can cause health problems, especially in the liver. A reference for evaluating liver function is blood test results for ALT, AST, ALP, and GGT activity. The objective of this study was to characterize the transaminase enzyme activity in horticultural farmers in the southern ring road area of Jambi city, based on the risk factors associated with pesticide exposure. This study employed a cross-sectional study approach in conjunction with a descriptive method. A total of thirty-four participants were involved, and blood samples were obtained from each for analysis in a lab. A photometer was utilized in the Medical Laboratory Technology department at Health Polytechnic of Jambi to measure the activity of liver enzymes. This study found some respondents who experienced increased enzyme activity, namely ALT: 8 people (23.5%), AST: 3 people (8.8%), ALP: 1 person (2.9%), and GGT: 1 person (2.9%). Based on the risk description observed, it was known that the intensity of pesticide exposure showed a significant increase only in ALT enzyme activity ($p=0.0048$), while adherence to mask-wearing increased ALT ($p=0.0018$) and GGT ($p=0.0134$). This study discovered that wearing a mask and the amount of pesticide exposure can increase enzyme activity, which may be a sign of liver impairment in the horticultural farmers under observation. It is anticipated that workers will pay greater attention to workplace safety by wearing masks and applying pesticides in the recommended dosages.

Keywords: Horticultural Farmers, Pesticide, Liver Enzyme Activity.

***Corresponding Author:**

James Perdinan Simanjuntak

Department of Environmental Health, Poltekkes Kemenkes Jambi, Jambi City, Jambi Province, Indonesia

Email: james.p.simanjuntak@poltekkesjambi.ac.id



©The Author(s) 2023. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.

1. INTRODUCTION

Indonesia is a productive country in the agricultural sector. Horticulture is one of the commodities produced by this sector. Considering that horticulture has characteristics that are easily damaged, managing pests and plant diseases has a significant impact on productivity (Amilia, Joy, and Sunardi., 2016). In reality, a large number of horticultural farmers continue to use excessive amounts of chemicals, such as pesticides, to enhance yields and product quality, which can have negative effects on health (Agustina & Norfai, 2018; Hassaan & El Nemr, 2020; Mrema et al., 2017).

According to data from the World Health Organization (WHO), there are between one and five million cases of poisoning among agricultural laborers annually, varying in severity and potentially affecting vital organs like the kidneys, lungs, or heart. An estimated 11,000 people die globally each year as a result of unintentional pesticide poisoning (Boedeker et al., 2020; Tostado & Bollmohr, 2022). Meanwhile, the Food and Drug Supervisory Agency reported that there were 344 cases of pesticide poisoning in Indonesia in 2019; 41.7% of those cases were associated with agricultural pesticides (BPOM, 2020)

Pesticides are generally absorbed by farmers through inhalation, digestion, or dermal, and distributed through the circulatory system to affect various organs, particularly the liver and kidneys involved in detoxification. The health of the farmers themselves may be adversely affected by this exposure condition (Boedeker et al., 2020; Jamal et al., 2015). ALT (alanine aminotransferase), also known as SGPT (serum glutamic Pyruvic transaminase), AST (aspartate aminotransferase), also known as SGOT (serum glutamic Oxaloacetic transaminase), GGT (gamma-glutamyl transaminase), and ALP (alkaline phosphatase), are manifestations of the enzymes connected to these organs and can be utilized to demonstrate the presence of liver dysfunction (Lala et al., 2022). Through measuring the activity of these enzymes in horticultural farmers in Jambi City's South Rim neighborhood who frequently come into contact with pesticides at work, a picture of the health of their livers was intended to be obtained.

The analysis was conducted using the following criteria: how long the research participants had been farmers, how frequently they implemented pesticides at work, and whether they employed masks, or personal protective equipment, to protect themselves from herbicide and insecticide exposure while spraying pesticides. Concerning farmers' attitudes and practices regarding the use of pesticides, as well as their elevated enzyme activities, which are considered to be an index marker of hepatotoxicity, these studies were conducted.

2. RESEARCH METHOD

Descriptive research utilizing a cross-sectional approach is the methodology employed. In Paal Merah, Jambi's Lingkar Selatan Village, sampling was conducted. The clinical chemistry laboratory of the Department of Medical Laboratory Technology at Health Polytechnic of Jambi is where the activity of the enzymes ALT, AST, ALP, and GGT was evaluated.

Horticultural farmers in Jambi's Lingkar Selatan Paalmerah Village who employed herbicides and insecticides contributed to the study's sample. Thirty-four farmers who belonged to Gapoktan Tani Makmur were the samples employed for this study. Purposive sampling was the method utilized for sampling, and 10 to 14 willing respondents from each group were selected to constitute the sample. The Health Polytechnic of Jambi's Research Ethics Committee granted ethical approval for this study, with approval number LB.02.06/2/353/2022.

Blood specimens from each subject were taken using a closed system with a yellow lid vacutainer tube filled with a gel separator.

To extract serum for testing, the frozen blood was centrifuged for five minutes at 3000 RPM. Using BioSystems reagent and the BTS-350 semi-automatic photometer, the kinetic method was used to measure the activity of the enzymes ALT, AST, ALP, and GGT.

After the enzyme activity was determined, the data was examined to discover how it related to the respondents' attributes, particularly those that were directly connected to their risk of pesticide exposure. The average value of enzyme activity and the frequency, or percentage, of instances of high enzyme activity detected are displayed in tabulations and graphs containing the data. Statistical different tests (ANOVA test and independent t-test) were performed using the Medcalc ver. 19.0.7 for Windows with a significance level of 0.05.

3. RESULTS AND DISCUSSION

Table 1. Characteristics of demographic and enzyme activity test results

Characteristics		Total	Percentage
Gender	Male	22	64.7%
	Female	12	35.3%
Age	Mean (\pm SD) *	41.6	(\pm 10.8)
	Adult	18	52.9%
	Pre-elderly	14	41.2%
Length of time as a farmer	Elderly	2	5.9%
	Mean (\pm SD) *	14.8	(\pm 8.7)
	<10 years	12	35.3%
Intensity of pesticide using	11-20 years	12	35.3%
	Very rarely	15	44.1%
	Rarely	13	38.2%
Mask-wearing adherence	Often	6	17.6%
	Very rarely	2	5.9%
	Rarely	9	26.5%
Alanine transaminase (ALT)	Always	23	67.6%
	Mean (\pm SD) *	30.3	(\pm 16.4)
	Normal	26	76.5%
Aspartate transaminase (AST)	High	8	23.5%
	Mean (\pm SD) *	23.6	(\pm 8.6)
	Normal	31	91.2%
Alkaline Phosphatase (ALP)	High	3	8.8%
	Mean (\pm SD) *	70.2	(\pm 25.1)
	Normal	33	97.1%
Gamma-glutamyl transferase (GGT)	High	1	2.9%
	Mean (\pm SD) *	22.7	(\pm 12.8)
	Normal	31	91.2%
	High	1	2.9%

* Mean and standard deviation (enzyme activity in IU/L; age & length of being a farmer in years)

The study's samples' clinical laboratory and demographic characteristics are displayed in Table 1. The Indonesian Ministry of Health categorized age characteristics into three groups: adults (19–44 years), pre-elderly (45–60 years), and elderly (>60 years) (Dahlan et al., 2018).

The duration of employment as a farmer was also classified into three groups according to the respondents' statements about the date of their initial horticultural employment. The predefined range in this study is every ten years. The frequency with which respondents administered pesticides while at work was divided into three categories: very rare (less than three times per month), rare (one to two times per week), and frequent (more than two times per week). Furthermore, data were also obtained on the level of respondents' compliance in wearing a good mask as a piece of personal protective equipment (PPE) when spraying pesticides in the categories of very rarely (never or only occasionally wearing a mask), rarely (sometimes wearing a mask), and often (almost always or always wearing a mask).

Table 1. illustrates that most of the respondents are male with the criteria of mature and pre-elderly age. The respondents' duration of farming is equitably distributed according to the three observed criteria. Based on the description of the intensity of spraying and the use of masks obtained, it appeared that the majority of respondents also had a relatively low risk of pesticide exposure. The average value of the test results for each of the observed laboratory parameters is also provided, and it is still within the normal range. The graph in Figure 1 below demonstrates how each enzyme activity test result was distributed among the 34 horticultural farmer respondents who were observed.




Figure 1. Boxplot graph of the results of measuring enzyme activity in horticultural farmers' liver function tests

The majority of the findings from the analysis of the four enzyme activities identified in this investigation decreased within the typical range for each parameter. Lala et al. (2022) provided the normal values for this study, which are as follows: GGT: 6-50 IU/L; AST: 5-30 IU/L; ALP: 30-120 IU/L; and ALT: 4-36 IU/L. Various frequencies of enzyme activity descending into the high category (exceeding the normal value) were observed for each enzyme test parameter. The largest proportion was identified in 8 respondents (23.5%) with high ALT activity test parameter results. ALP had one respondent (2.9%), GGT had one respondent (2.9%), and AST had three (8.8%) respondents. In the meantime, these were the other three parameters.

Based on the results of the demographic features of the study participants, the percentage of high-activity events in the four enzymes that were observed was subsequently examined (figure 2). The group of respondents who did not wear protective equipment (PPE) included individuals with high ALT and AST activities. In contrast to variations from other characteristics, respondents with the characteristic of the intensity of pesticide spraying that

was included in the frequent category had the highest percentage of high ALP and GGT enzyme activities. Based on the characteristics of each respondent group examined in this study, overall ALT has the highest percentage when compared to other parameters.

Toxic chemical compounds can enter the human body in several ways, encompassing absorption through the skin, orally, or inhalation, either intentionally or unintentionally. These dangerous materials are a potential workplace risk for many different professions worldwide. The detrimental effects of pesticide exposure on liver disorders, which are in charge of neutralizing dangerous substances that enter the body, have been documented in a number of long-running studies (Colombo et al., 2019; Luo et al., 2005; Malaguarnera, 2012; Redlich, 1988). Prospective cohort studies have also reported on discussions regarding different pesticide types that affect human health. These studies state that, despite not having a high and consistent potential, this hazardous material is predicted to be one of the triggers of cancer in a person, involving bladder and liver cancer (Koutros et al., 2016; Rapisarda et al., 2016; VoPham et al., 2017).

Studies on various types of farmers have also reported pesticide exposure. The liver eventually suffers chronic damage from the accumulation of this exposure (Melaram, 2021; Damalas & Koutroubas, 2016). Based on the results of the study's analysis of the respondents' demographic features, the percentage of high activity occurrence in each of the four enzymes that were identified was further examined (Figure 2). The "very rarely" category of respondents' mask-wearing compliance displayed the highest percentage (100%) for the ALT and AST parameters. In contrast to other risk variations or characteristics, the highest percentage (50%) of respondents with pesticide spraying intensity in the "frequent" category had high activity of ALP and GGT enzymes.




Figure 2. Graph of the percentage of high enzyme activity based on the characteristics of horticultural farmers.

Statistical analysis was administered to show differences in the results of the four parameters of measurement of enzyme activity based on each variable group of respondent characteristics. There were no discernible differences between the respondents' age, gender, or length of time employed as farmers for any of the enzyme activities. Stated differently, there was no observed increase in enzyme activity among the respondents as a result of these three variables. There was a difference in the ALT enzyme activity measurements according to the degree of pesticide exposure ($p=0.0048$) and farmers' adherence to wearing masks when spraying ($p=0.0018$). Furthermore, this significance was also demonstrated by the GGT enzyme activity, but only in the variable mask use ($p=0.0134$). In contrast, AST and ALP activities did not provide a significant difference in each of the variations of these two variables.

Table 2. Analysis of the characteristics of farmers on the results of enzyme activity tests.

Characteristic	ALT			AST			ALP			GGT		
	N	H	Mean	N	H	Mean	N	H	Mean	N	T	Mean
Gender												
Male	18	4	27.3	20	2	23.3	21	1	69.5	21	1	20.7
Female	8	4	35.8	11	1	24.2	12	0	72.1	12	0	26.4
p-value			0.125			0.3295			0.705			0.1251
Age group												
Adult	14	4	29.7	17	1	21.7	18	0	73.3	17	1	21.5
Pre-elderly	10	4	32.4	13	1	24.9	13	1	66.1	14	0	23.9
Elderly	2	0	21	1	1	32	2	0	74.5	2	0	25.5
p-value			0.6511			0.2173			0.7307			0.8354
Length of time as a farmer												
<10 years	10	2	29	12	0	20.7	12	0	74.1	12	0	20.2
11-20 years	7	5	35	11	1	23.3	11	1	69.5	11	1	26.3
>20 years	9	1	26.2	8	2	27.4	10	0	67.1	10	0	25.1
p-value			0.4456			0.1917			0.819			0.4822
Intensity of pesticide-using												
Very rarely	15	0	21.4	14	1	20.7	15	0	67.3	15	0	17.9
Rarely	8	5	33.9	12	1	25.7	13	0	73.5	13	0	25.9
Often	3	3	44.7	5	1	27.5	5	1	71.9	5	1	27.8
p-value			0.0048			0.1138			0.8206			0.1462
			**									
Mask-wearing adherence												
Very rarely	0	2	25.3	0	2	21.5	2	0	69.4	2	0	19.2
Rarely	5	4	35.8	8	1	28.2	9	0	69.4	8	1	27.1
Always	21	2	63	23	0	26.5	22	1	87.2	23	0	43.5
p-value			0.0018			0.1252			0.6496			0.0134
			**									*

Note: N = number of respondents with normal enzyme activity; H = the number of respondents with high enzyme activity; Mean = average value of enzyme activity (IU/L); P-value is significant at the level of: * = 0.05 & ** = 0.01

In comparison to other parameters, overall ALT is also the one that frequently demonstrates the highest percentage, particularly when it comes to respondent characteristics associated with pesticide exposure risk factors. Although pesticide poisoning also increases other enzyme activities present in farmers' blood, this is very consistent with the phenomenon

found in several previous studies in several countries that showed an increase in the same enzyme in cases of pesticide poisoning (Damalas & Koutroubas, 2016; Freire et al., 2015; Jamal et al., 2015). When Lozano-Paniagua et al., (2021) observed greenhouse farmers in Spain who were exposed to pesticides, they discovered that workers who had high exposure periods had higher levels of ALT and ALP. Bernieri et al., (2021) observed that although AST enzyme activity increased significantly following high pesticide exposure in Brazilian farmers, ALT and GGT did not increase. In the past, Manfo et al., (2020) used a case-control methodology to track the prevalence of kidney and liver diseases among farmers in Buea, Cameroon. The study revealed that exposure to pesticides increased the activity of the ALT enzyme significantly, but not that of the AST, creatinine, or uric acid. Nevertheless, because the majority of these enzymes are produced in the liver (hepatic cells), ALT is recognized as the most specific enzyme to indicate impaired liver function when compared to other enzymes (Aulbach & Amuzie, 2017).

Freire et al., (2015) who observed Brazilians who were heavily exposed to organochlorine pesticides also gathered that there was an increase in liver enzyme activity due to these compounds. Moreover, other parameters such as bilirubin, which is also the main marker of liver function abnormalities, were successfully evidenced to possess a significant relationship with beta-hexachlorocyclohexane levels unveiled in the blood of respondents. Furthermore, from the hematological aspect, eosinophilia was also reported, as well as decreased hemoglobin levels and erythrocyte counts in the blood.

Several factors, including age, length of employment, use of personal protective equipment (PPE), pesticide dosage, wind direction during spraying, and frequency of spraying, can affect how severe liver function disorders caused by pesticide exposure can be. Because farmers are more susceptible to come into direct contact with pesticides, including both herbicides and insecticides, the length of time they work can raise their risk of pesticide poisoning (Andarini & Rosanti, 2018). Farmers frequently disregard the importance of wearing personal protective equipment (PPE), such as masks, even though it can lessen the negative health effects of pesticide exposure. Numerous studies conducted on lab animals have demonstrated the hepatotoxic effects of pesticides (Nieradko-Iwanicka & Borzęcki, 2015; Rizzati et al., 2016). Meanwhile, Yokoyama et al., (2019) asserted that exposure to pesticides provided to animals tested in his study did not cause an increase in ALP enzyme activity. This is similar to the facts uncovered in this study on ALP enzyme activity which also did not perform any significant difference from each risk group observed.

Bayili et al., (2020) performed a longitudinal study on 113 farmers in two phases: the first phase occurred during the harvest period, and the second phase was conducted six months after the harvest period ended. The second phase's ALT and AST activity significantly decreased ($p < 0.01$), according to the results. Although ALP's average value decreased as well, there was no discernible change. On the other hand, GGT activity rose from 40.5 ± 30.68 U/L in the first phase to 63.0 ± 67.07 U/L in the second phase ($p < 0.0001$). This demonstrates how heavily farmers' exposure to pesticides during their work affects abnormalities in liver function; the less intensely farmers are exposed, the lower the results of the measurement of liver enzyme activity that exceeds normal values.

In a study conducted in Indonesia, Sukmayanti et al., (2020) concluded that gender and age affect ALT and cholinesterase enzyme activity. Male farmers in Bali's Tabanan district had higher levels of enzyme activity than female farmers, according to observations made of them growing vegetables. This contrasts with the phenomenon uncovered when ALT activity was measured in farmers in Jambi City's South Ring, where nearly all of the enzymes discovered had mean values that were higher in the female group than the male group. In the same way, the researchers identified no statistically significant differences in the age group of farmers in

this study, even though the observed enzyme activity values in the South Ring of Jambi City tended to increase with farmer age.

Meanwhile, research conducted by Tsani et al., (2017) revealed that 29 out of 43 farmers (67.4%) in Sumberejo village, Magelang had impaired liver function. The study ultimately concluded that a working period can have an impact on liver function impairment and lead to an accumulation of pesticides in the body. It was determined that there is little correlation between PPE use and the prevalence of liver function disorders. This is conceivable because there was only one farmer respondent in the study who fully utilized PPE. The results of a study on farmers in Jambi City's South Ring demonstrate a lower percentage—just 9 out of 34 farmers, or 26.5 percent. Since the length of service as a farmer generated no statistically significant results in this study, it can be assumed that it is not a significant factor in determining the onset of liver function disorders. The high intensity of pesticide use and not followed by the use of good PPE is proven to cause the accumulation of pesticides in the body to increase and cause impaired liver function.

4. CONCLUSION

Within the southern ring road of Jambi City, nine out of the horticultural farmers who used pesticides (26.5%) had high enzyme measurement results, according to this study. The most frequently encountered enzyme with elevated activity above normal values was alanine aminotransferase (ALT). This study has demonstrated that the intensity of pesticide spraying and mask use by horticultural farmers are risk factors that may lead to elevated liver enzyme activity, particularly ALT and GGT.

REFERENCES

- Agustina, N., & Norfai, N. (2018). Paparan Pestisida terhadap Kejadian Anemia pada Petani Hortikultura. *Majalah Kedokteran Bandung*, 50(4), 215–221. <https://doi.org/10.15395/mkb.v50n4.1398>
- Amilia, E., Joy, B., & Sunardi, S. (2016). Residu Pestisida pada Tanaman Hortikultura (Studi Kasus di Desa Cihanjuang Rahayu Kecamatan Parongpong Kabupaten Bandung Barat). *Agrikultura*, 27(1), 23-29. <https://doi.org/10.24198/agrikultura.v27i1.8473>
- Andarini, Y. D., & Rosanti, E. (2018). Kajian toksisitas pestisida berdasarkan masa kerja dan personal hygiene pada petani hortikultura di Desa Demangan. *An-Nadaa: Jurnal Kesehatan Masyarakat (e-Journal)*, 5(2), 82-89. <https://doi.org/10.31602/ann.v5i2.1655>
- Aulbach, A. D., & Amuzie, C. J. (2017). Biomarkers in Nonclinical Drug Development. In *A Comprehensive Guide to Toxicology in Nonclinical Drug Development*, pp. 447–471. Elsevier. <https://doi.org/10.1016/B978-0-12-803620-4.00017-7>
- Bayili, B., Da, O., Ilboudo, S., Ouedraogo, R., Coulibaly, V. P., Bationo, J. F., ... & Ouedraogo, G. A. (2020). Study of biochemical parameters in farmers exposed to pesticides used in cotton growing around the Bala hippopotamus pond. *GSC Biological and Pharmaceutical Sciences*, 13(1), 111-122. <https://doi.org/10.30574/gscbps.2020.13.1.0323>
- Bernieri, T., Rodrigues, D., Randon Barbosa, I., Perassolo, M. S., Grolli Ardenghi, P., & Basso da Silva, L. (2021). Effect of pesticide exposure on total antioxidant capacity and biochemical parameters in Brazilian soybean farmers. *Drug and Chemical Toxicology*, 44(2), 170–176. <https://doi.org/10.1080/01480545.2019.1566353>
- Boedeker, W., Watts, M., Clausing, P., & Marquez, E. (2020). The global distribution of acute unintentional pesticide poisoning: Estimations based on a systematic review. *BMC Public Health*, 20(1), 1875. <https://doi.org/10.1186/s12889-020-09939-0>
- BPOM. (2020). *Laporan Tahunan Pusat Data dan Informasi Obat dan Makanan tahun 2019*. Badan Pengawasan Obat dan Makanan.

- Colombo, M., La Vecchia, C., Lotti, M., Lucena, M. I., Stove, C., Paradis, V., & Newsome, P. (2019). EASL Clinical Practice Guideline: Occupational liver diseases. *Journal of Hepatology*, 71(5), 1022–1037. <https://doi.org/10.1016/j.jhep.2019.08.008>
- Dahlan, A. K., Umrah, A. S., & Abeng, T. (2018). Kesehatan Lansia: Kajian Teori Gerontologi dan Pendekatan Asuhan pada Lansia. Malang: Intimedia.
- Damalas, C., & Koutroubas, S. (2016). Farmers' Exposure to Pesticides: Toxicity Types and Ways of Prevention. *Toxics*, 4(1), 1. <https://doi.org/10.3390/toxics4010001>
- Freire, C., Koifman, R. J., & Koifman, S. (2015). Hematological and Hepatic Alterations in Brazilian Population Heavily Exposed to Organochlorine Pesticides. *Journal of Toxicology and Environmental Health, Part A*, 78(8), 534–548. <https://doi.org/10.1080/15287394.2014.999396>
- Hassaan, M. A., & El Nemr, A. (2020). Pesticides pollution: Classifications, human health impact, extraction and treatment techniques. *The Egyptian Journal of Aquatic Research*, 46(3), 207–220. <https://doi.org/10.1016/j.ejar.2020.08.007>
- Jamal, F., Haque, Q. S., Singh, S., & Arshad, M. (2015). The Influence of Pesticides on Hepatic and Renal Functions in Occupational Sprayers of Rural Malihabad, Lucknow (India). *Toxicology: Open Access*, 1(01). <https://doi.org/10.4172/2476-2067.1000106>
- Koutros, S., Silverman, D. T., Alavanja, M. C., Andreotti, G., Lerro, C. C., Heltshe, S., Lynch, C. F., Sandler, D. P., Blair, A., & Beane Freeman, L. E. (2016). Occupational exposure to pesticides and bladder cancer risk. *International Journal of Epidemiology*, 45(3), 792–805. <https://doi.org/10.1093/ije/dyv195>
- Lala, V., Zubair, M., & Minter, D. A. (2022). *Liver Function Tests*. StatPearls. Treasure Island (FL). Retrieved from <https://pubmed.ncbi.nlm.nih.gov/29494096/>
- Lozano-Paniagua, D., Parrón, T., Alarcón, R., Requena, M., López-Guarnido, O., Lacasaña, M., & Hernández, A. F. (2021). Evaluation of conventional and non-conventional biomarkers of liver toxicity in greenhouse workers occupationally exposed to pesticides. *Food and Chemical Toxicology*, 151, 112127. <https://doi.org/10.1016/j.fct.2021.112127>
- Luo, J.-C., Cheng, T.-J., Kuo, H.-W., & Chang, M. J. W. (2005). Abnormal liver function associated with occupational exposure to dimethylformamide and glutathione S -transferase polymorphisms. *Biomarkers*, 10(6), 464–474. <https://doi.org/10.1080/13547500500333648>
- Malaguarnera, G. (2012). Toxic hepatitis in occupational exposure to solvents. *World Journal of Gastroenterology*, 18(22), 2756. <https://doi.org/10.3748/wjg.v18.i22.2756>
- Manfo, F. P. T., Mboe, S. A., Nantia, E. A., Ngoula, F., Telefo, P. B., Moundipa, P. F., & Cho-Ngwa, F. (2020). Evaluation of the Effects of Agro Pesticides Use on Liver and Kidney Function in Farmers from Buea, Cameroon. *Journal of Toxicology*, 2020, 1–10. <https://doi.org/10.1155/2020/2305764>
- Melaram, R. (2021). Environmental Risk Factors Implicated in Liver Disease: A Mini-Review. *Frontiers in Public Health*, 9, 683719. <https://doi.org/10.3389/fpubh.2021.683719>
- Mrema, E. J., Ngowi, A. V., Kishinhi, S. S., & Mamuya, S. H. (2017). Pesticide Exposure and Health Problems Among Female Horticulture Workers in Tanzania. *Environmental Health Insights*, 11, 117863021771523. <https://doi.org/10.1177/1178630217715237>
- Nieradko-Iwanicka, B., & Borzęcki, A. (2015). Effect of 28-Day Exposure to Fenpropathrin on the Activities of Serum Alanine Transaminase and Liver Antioxidant Enzymes in Mice. *Bulletin of the Veterinary Institute in Pulawy*, 59(1), 165–169. <https://doi.org/10.1515/bvip-2015-0025>
- Rapisarda, V., Loreto, C., Malaguarnera, M., Ardiri, A., Proiti, M., Rigano, G., Frazzetto, E., Ruggeri, M. I., Malaguarnera, G., Bertino, N., Malaguarnera, M., Catania, V. E., Di Carlo, I., Toro, A., Bertino, E., Mangano, D., & Bertino, G. (2016). Hepatocellular

- carcinoma and the risk of occupational exposure. *World Journal of Hepatology*, 8(13), 573. <https://doi.org/10.4254/wjh.v8.i13.573>
- Redlich, C. A. (1988). Liver Disease Associated with Occupational Exposure to the Solvent Dimethylformamide. *Annals of Internal Medicine*, 108(5), 680. <https://doi.org/10.7326/0003-4819-108-5-680>
- Rizzati, V., Briand, O., Guillou, H., & Gamet-Payraastre, L. (2016). Effects of pesticide mixtures in human and animal models: An update of the recent literature. *Chemico-Biological Interactions*, 254, 231–246. <https://doi.org/10.1016/j.cbi.2016.06.003>
- Sukmayanti, N. L. P. A., Artini, N. P. R., & Yanti, N. P. W. (2020). Analisis Kadar SGPT (Serum Glutamic Pyruvic Transaminase) Dan Kholinesterase Pada Petani Sayur Di Desa Riang Gede, Kecamatan Penebel, Kabupaten Tabanan. *THE JOURNAL OF MUHAMMADIYAH MEDICAL LABORATORY TECHNOLOGIST*, 3(2), 25. <https://doi.org/10.30651/jmlt.v3i2.5841>
- Tostado, L., & Bollmohr, S. (2022). Facts and figures about toxic chemicals in agriculture 2022. *Creative Commons, 1st edition*. Retrieved from <https://eu.boell.org/PesticideAtlas>
- Tsani, R. A., Setiani, O., & Dewanti, N. A. Y. (2017). Hubungan Riwayat pajanan pestisida dengan gangguan fungsi hati pada petani di desa sumberejo kecamatan ngablak kabupaten magelang. *Jurnal Kesehatan Masyarakat*, 5(3), 411-419. <https://doi.org/10.14710/jkm.v5i3.17258>
- VoPham, T., Bertrand, K. A., Hart, J. E., Laden, F., Brooks, M. M., Yuan, J.-M., Talbott, E. O., Ruddell, D., Chang, C.-C. H., & Weissfeld, J. L. (2017). Pesticide exposure and liver cancer: A review. *Cancer Causes & Control*, 28(3), 177–190. <https://doi.org/10.1007/s10552-017-0854-6>
- Yokoyama, Y., Ono, A., Yoshida, M., Matsumoto, K., & Saito, M. (2019). Toxicological significance of increased serum alkaline phosphatase activity in dog studies of pesticides: Analysis of toxicological data evaluated in Japan. *Regulatory Toxicology and Pharmacology*, 109, 104482. <https://doi.org/10.1016/j.yrtph.2019.104482>